



common name: black soldier fly

scientific name: *Hermetia illucens* (Linnaeus) (Insecta: Diptera: Stratiomyidae)

[Introduction](#) - [Synonymy](#) - [Distribution](#) - [Description and Life Cycle](#) - [Economic Importance](#) - [Selected References](#)

Introduction ([Back to Top](#))

The black soldier fly, *Hermetia illucens* (Linnaeus), is a sleek looking fly that many confuse with a wasp. However, like most flies, the black soldier flies only have two wings (wasps have four) and does not possess a stinger. Although the loud buzzing they create when flying is enough to concern many people, adult soldier flies pose no danger.



Figure 1. Adult black soldier fly, *Hermetia illucens* (Linnaeus). Photograph by Lyle J. Buss, University of Florida.

The black soldier fly is often associated with the outdoors and livestock, usually around decaying organic matter such as animal waste or plant material. Since black soldier fly larvae consume decaying matter, they have been used to reduce animal manure in commercial swine and poultry facilities (Newton 2005). Although they are not known as a disease vector, adult soldier flies are a potential mechanical vector of various pathogens. A more likely negative interaction would be accidental ingestion of black soldier fly larvae by animals or humans. (Goddard 2003).

Synonymy ([Back to Top](#))

Musca illucens Linnaeus, 1758

Musca leucopa Linnaeus, 1767

Hermetia rufiventris Fabricius, 1805

Hermetia pellucens Macquart, 1834

Hermetia nigrifacies Bigot, 1879

Hermetia mucens Riley & Howard, 1889

Hermetia illucens var. *nigritibia* Enderlein, 1914

Hermetia illucens Copello, 1926

List generated from (ITIS 2009)

Distribution [\(Back to Top\)](#)

The family Stratiomyidae comprises 260 known species in North America (Triplehorn 2005). In the southeastern United States, the black soldier fly is abundant during late spring and early fall, and has three generations per year in Georgia (Tomberlin et al. 2002). While common in the continental United States, this fly is found throughout the Western Hemisphere.

In natural breeding sites (i.e. carrion) black soldier flies lay their eggs in moist organic material. Black soldier flies frequent agricultural settings because the large amounts of organic waste left by livestock offer abundant sites that meet their reproductive needs. In areas where natural habitats are removed (urbanized areas) the black soldier fly will lay eggs in dumpsters or compost, which provide similar odors and nutritional needs to naturally occurring organic matter. This is especially true for areas with poor sanitation.

Description and Life Cycle [\(Back to Top\)](#)

Adults: Members of the soldier fly family Stratiomyidae can range in color from yellow, green, black or blue, with some having a metallic appearance. Many are mimics of other flying insects, such as bees and wasps. Black soldier fly adults have a wasp-like appearance and are black or blue in color. Soldier flies also have two translucent "windows" located on the first abdominal segment. Adults range from 15 to 20 mm in length. (Sheppard et al. 2002). The adult's antennae are elongated with three segments, and legs have white coloration near the end of each leg.



Figure 2. Lateral view of an adult black soldier fly, *Hermetia illucens* (Linnaeus). Notice the two translucent "windows" located on the first abdominal segment, partially obscured in this image by the rear leg. Photograph by Bianca Diclaro, University of Florida.

Two days after adult emergence from the pupal case, mating can occur. A male black soldier fly intercepts a passing female in mid-air and they descend *in copula* (Tomberlin and Sheppard 2001). Male soldier flies utilize lekking sites, where they await female soldier flies. These sites are defended against other male soldier flies. When a male intrudes upon the territory of a resting male, the resting male seizes the intruder. After a brief descent, the invading male will retreat.

Eggs: The female black soldier fly deposits a mass of about 500 eggs in cracks and crevices near or in decaying matter such as dung, carrion, garbage, and other organic waste. The eggs hatch into larvae in

about four days. Each oval shaped egg is about 1 mm in length, and pale yellow or creamy white in color (NCIPMI 1998).

Larvae: The larvae can reach 27 mm in length and 6 mm in width. They are a dull, whitish color with a small, projecting head containing chewing mouthparts. Larvae pass through six instars and require approximately 14 days to complete development (Hall and Gerhardt 2002). During larval development, black soldier fly larvae are insatiable feeders. As adults they do not need to feed and rely on the fats stored from the larval stage (Newton et al. 2005).



Figure 3. Dorsal (left) and ventral (right) views of sixth instar larvae of the black soldier fly, *Hermetia illucens* (Linnaeus). Photograph by Bianca Diclaro, University of Florida.

Pupae: Before pupation, the sixth instar larvae disperse from the feeding site to dry sheltered areas, such as ground vegetation, to initiate pupation. The exoskeleton (skin) darkens and a pupa develops within. Pupation requires about two weeks (Hall and Gerhardt 2002).



Figure 4. Pupa of the black soldier fly, *Hermetia illucens* (Linnaeus). Photograph by James Castner, University of Florida.

Economic Importance [\(Back to Top\)](#)

The adult black soldier fly is not usually considered a pest (Newton et al. 2005). Because the larvae have been shown to be effective manure recyclers, a "Black Soldier Fly Manure Management System" has been proposed to not only reduce livestock waste, but also generate a food source for fish and other animals. In a program outlined in Newton et al. (2005) swine manure was fed to black soldier fly larvae, which greatly reduced the waste material. The manure was transferred into a basin containing black soldier fly larvae. As the larvae developed they reduced the manure by 50%. Approximately 45,000 larvae will consume 24 kg of swine manure in 14 days. As the larvae mature they crawl out of the basin, thereby self-harvesting themselves, and are subsequently available as livestock feed. In addition to being a good source of oil and protein for animal feed, black soldier fly larvae have the potential of improving organic waste into a rich fertilizer.



Figure 5. Larvae of the black soldier fly, *Hermetia illucens* (Linnaeus), in compost.

Selected References [\(Back to Top\)](#)

- Drees BM, Jackman J. (October 2008). Soldier fly. *A Field Guide to Common Texas Insects*. <http://insects.tamu.edu/fieldguide/cimg226.html> (14 July 2009).
- Goddard, J. 2003. Physician's Guide to Arthropods of Medical Importance, 4th Edition. CRC Press LLC. Boca Raton, Florida.
- Hall DC, Gerhardt RR. 2002. Flies (Diptera), pp 127-161. *In* Mullen G, Durden L. (editors). Medical and Veterinary Entomology. Academic Press. San Diego, California.
- ITIS. (2009). *Hermetia illucens* Linnaeus. *Integrated Taxonomic Information System*. http://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=130298 (14 July 2009).
- NCIPMI. (1998). Insect and related pests of man and animals. *North Carolina Integrated Pest Management Information*. http://ipm.ncsu.edu/AG369/notes/black_soldier_fly.html (14 July 2009).
- Newton L, Sheppard C, Watson DW, Burtle G, Dove, R. (June 2005). Using the black soldier fly, *Hermetia illucens*, as a value-added tool for the management of swine manure. *Waste Management Programs*. North Carolina State University. http://www.cals.ncsu.edu/waste_mgt/smithfield_projects/phase2report05/cd_web%20files/A2.pdf (14 July 2009).
- Sheppard DC, Tomberlin JK, Joyce JA, Kiser BC, Sumner SM. 2002. Rearing methods for the black soldier fly (Diptera: Stratiomyidae). *Journal of Medical Entomology* 39: 695-698.
- Tomberlin JK, Sheppard DC. 2001. Lekking behavior of the black soldier fly (Diptera: Stratiomyidae). *Florida Entomologist* 84: 729-730.
- Tomberlin JK, Sheppard DC, Joyce JA. 2002. Selected life history traits of black soldier flies (Diptera: Stratiomyidae) reared on three artificial diets. *Annals of the Entomological Society of America* 95: 379-386.
- Triplehorn CA, Johnson NF. 2005. Borror and Delong's Introduction to the Study of Insects, 7th Edition. Thomson Brooks/Cole. Belmont, California. 864 pp.



Authors: Joseph W. Diclaro II and [Phillip E. Kaufman](#), University of Florida
Photographs: Bianca Diclaro, [Lyle J. Buss](#), and James Castner, University of Florida
Web Design: Don Wasik, Jane Medley
Publication Number: EENY-461
Publication Date: July 2009. Reviewed: April 2018.

[Copyright University of Florida](#) ~ [An Equal Opportunity Institution](#)
Featured Creatures Editor and Coordinator: [Jennifer L. Gillett-Kaufman](#), University of Florida



Florida Department of
Agriculture and Consumer Services
Adam H. Putnam, Commissioner

